

**Track Changes**  
**from Chapter 3, Section H V1.08**  
**to Chapter 3, Section H V1.10**

Chapter	Section	Page	Change
3	H0100	H-2	— In men, eCondom catheters (males), and in females, external urinary pouches (females), are commonly often used intermittently or at night only. This use should be; these should be coded as external catheters.
3	H0100	H-2	<div> <p><b>DEFINITIONS</b></p> <p><b>EXTERNAL CATHETER</b>  Device attached to the shaft of the penis like a condom for males or a receptacle pouch that fits around the labia majora majora for females and connected to a drainage bag.</p> </div>
3	H0200	H-4	<i>The look-back period for this item is to since the most recent admission/entry or reentry admission or since urinary assessment, the most recent prior assessment, or to when incontinence was first noted within the facility.</i>
3	H0200	H-5	<ul style="list-style-type: none"> <li>Code 1, yes: for residents who underwent a trial of an individualized, resident- centered toileting program at least once since admission/readmission, prior assessment prior assessment, or when urinary incontinence was first noted.</li> </ul>
3	H0300	H-7	<div> <p><b>DEFINITIONS</b></p> <p><b>URINARY INCONTINENCE</b>  The involuntary loss of urine.</p> <p><b>CONTINENCE</b>  Any void into a commode, urinal, or bedpan that occurs voluntarily, or voluntarily, or as the result of prompted toileting, assisted toileting, or scheduled toileting.</p> </div>
3	H0600	H-12	— side effects of medications.